arXiv:1207.6765v1 [math.CO] 29 Jul 2012

The Nullity of Bicyclic Signed Graphs*

Yi-Zheng Fan[†], Wen-Xue Du, Chun-Long Dong

School of Mathematical Sciences, Anhui University, Hefei 230601, P. R. China

Abstract: Let Γ be a signed graph and let $A(\Gamma)$ be the adjacency matrix of Γ. The nullity of Γ is the multiplicity of eigenvalue zero in the spectrum of $A(\Gamma)$. In this paper we characterize the signed graphs of order n with nullity $n-2$ or $n-3$, and introduce a graph transformation which preserves the nullity. As an application we determine the unbalanced bicyclic signed graphs of order n with nullity $n-3$ or $n-4$, and signed bicyclic signed graphs (including simple bicyclic graphs) of order n with nullity $n-5$.

Keywords: Signed graph; nullity; bicyclic graphs

MR(2000) Subject Classification: 05C50

1 Introduction

A *signed graph* is a graph with a sign attached to each of its edges. Formally, a signed graph $\Gamma = (G, \sigma)$ consists of a simple graph $G = (V, E)$, referred to as its underlying graph, and a mapping $\sigma : E \to \{+, -\}$, the edge labeling. To avoid confusion, we also write $V(\Gamma)$ or $V(G)$ instead of V, $E(G)$ instead of E, and $E(\Gamma) = E^\sigma$. The underlying graph G of Γ is sometimes denoted by ${}^u\Gamma$.

The *adjacency matrix* of Γ is $A(\Gamma) = (a^\sigma_{ij})$ with $a^\sigma_{ij} = \sigma(v_i v_j) a_{ij}$, where (a_{ij}) is the adjacency matrix of the underlying graph G. In the case of $\sigma = +$, which is an all-positive edge labeling, $A(G, +)$ is exactly the classical adjacency matrix of G. So a *simple graph* is always assumed as a signed graphs with all edges positive. The *nullity of a signed graph* Γ is defined as the multiplicity of the eigenvalue zero in the spectrum of $A(\Gamma)$, and is denoted by $\eta(\Gamma)$. The *rank* of Γ is referred to the rank of $A(\Gamma)$, and is denoted by $r(\Gamma)$. Surely $\eta(\Gamma) + r(\Gamma) = n$ if Γ has n vertices.

Let C be a cycle of Γ. The *sign* of C, denoted by $\operatorname{sgn}(C)$, and is defined as $\operatorname{sgn}(C) = \Pi_{e \in C} \sigma(e)$. The cycle C is said to be *positive* or *negative* if $\operatorname{sgn}(C) = +$ or $\operatorname{sgn}(C) = -$. A signed graph is said to be *balanced* if all its cycles are positive, or equivalently, all cycles have an even number of negative edges; otherwise it is called *unbalanced*.

Suppose $\theta : V(G) \to \{+, -\}$ is any sign function. Switching Γ by θ means forming a new signed graph $\Gamma^\theta = (G, \sigma^\theta)$ whose underlying graph is the same as G, but whose sign function is defined on an edge uv by $\sigma^\theta(uv) = \theta(u)\sigma(uv)\theta(v)$. Note that switching does not change the signs or balanceness of the cycles of Γ. If we define a (diagonal) signature matrix D^θ with $d_v = \theta(v)$ for each $v \in V(G)$, then $A(\Gamma^\theta) = D^\theta A(\Gamma) D^\theta$. Two graphs Γ_1, Γ_2 are called *switching equivalent*, denoted by $\Gamma_1 \sim \Gamma_2$, if there exists a switching function θ such that $\Gamma_2 = \Gamma_1^\theta$, or equivalently $A(\Gamma_2) = D^\theta A(\Gamma_1) D^\theta$.

Theorem 1.1. [20] *Let Γ be a signed graph. Then Γ is balanced if and only if $\Gamma = (G, \sigma) \sim (G, +)$.*

---

*Supported by National Natural Science Foundation of China (11071002, 11126178), Program for New Century Excellent Talents in University, Key Project of Chinese Ministry of Education (210091), Specialized Research Fund for the Doctoral Program of Higher Education (20103401110002), Science and Technological Fund of Anhui Province for Outstanding Youth (10040606Y33), Project of Educational Department of Anhui Province (KJ2011A019), Scientific Research Fund for Fostering Distinguished Young Scholars of Anhui University(KJJQ1001), Academic Innovation Team of Anhui University Project (KJTD001B).

[†] Corresponding author. E-mail addresses: fanyz@ahu.edu.cn (Y.-Z. Fan), wenxuedu@gmail.com (W.-X. Du), 786189560@qq.com (C.-L. Dong).

Recently the authors [12] investigate the nullity of signed graphs, and determine the unicyclic signed graphs with nullity $n-2$, $n-3$, $n-4$, or $n-5$. In this paper we characterize the signed graphs of order n with nullity $n-2$ or $n-3$, and introduce a graph transformation which preserves the nullity. As an application we determine the unbalanced bicyclic signed graphs of order n with nullity $n-3$ or $n-4$, and signed bicyclic signed graphs (including simple bicyclic graphs) of order n with nullity $n-5$. We hope this work would be helpful to characterize the structure of signed graphs from the view of nullity.

Finally we note some work on the nullity of simple graphs. For a simple graph G of order n containing at least one edge, $0 \leq \eta(G) \leq n-2$. Cheng and Liu [3] characterize the simple graphs of order n with nullity $n-2$ or $n-3$. Cheng, Huang and Yeh [4, 5] characterize the simple graphs of order n with nullity $n-4$ or $n-5$. The characterization of simple graphs of order n with nullity $n-6$ or more is still open. Much work is devoted to the nullity of special classes of simple graphs; see [11, 13, 15, 16, 17, 18, 22, 25, 26, 27, 29, 32, 34]. The work on the nullity of simple graphs were found to be useful in chemistry to investigate the stability of the alternant hydrocarbons ([6],[28]), and is also of interest in mathematics itself, as it is closely related to the minimum rank problem of symmetric matrices whose patterns are described by graphs [9].

2 Nullity of signed graphs

Some notations are listed below. Let Γ be a signed graph. Let v be a vertex of Γ and let $U \subset V(\Gamma)$. Denote by $N(v)$ the neighborhood of v in Γ, $d(v)$ the degree of v (i.e. the cardinality of $N(v)$). Denote by $\Gamma[U]$ the induced subgraph of Γ on the vertices of U including the signs of edges. Denote by P_r, C_r, K_r respectively a path, a cycle and a complete graph, all of which are simple graphs of order r. We use the notation $G_1 + G_2$ to denote the union of two vertex-disjoint simple graphs G_1, G_2.

Lemma 2.1. [12] (i) *If T is an acyclic signed graph or a signed tree of order n. Then $\eta(T) = n - 2\mu(T)$.*

(ii) *If C_n is a balanced cycle, then $\eta(C_n) = 2$ if $n \equiv 0(\bmod 4)$ and $\eta(C_n) = 0$ otherwise.*

(iii) *If C_n is an unbalanced signed cycle, then $\eta(C_n) = 2$ if $n \equiv 2(\bmod 4)$, and $\eta(C_n) = 0$ otherwise.*

If a signed graph Γ contains at least one edge, then $\eta(\Gamma) \leq n-2$. By adopting the technique in [3], We first characterize the signed graphs of order n with nullity $n-2$ or $n-3$.

Lemma 2.2. *Suppose that Γ is a signed graph of order n and Γ has no isolated vertex. Let x be an arbitrary vertex in Γ. Let $Y = N(x)$ and $X = V(\Gamma) - Y$. If $r(\Gamma) \leq 3$, then*

(i) *No two vertices in X are adjacent.*

(ii) *Each vertex from X and each vertex from Y are adjacent.*

Proof. Let x_1, x_2 be two vertices in X. If one of them is x, surely the assertion (i) holds. Now assume x_1 and x_2 are adjacent and neither of them is x. Since Γ contains no isolated vertices, Y is not empty. Select any vertex y in Y. Then ${}^u\Gamma[x_1, x_2, y]$ is isomorphic to $P_2 + K_1$, P_3 or K_3, and hence ${}^u\Gamma[x, x_1, x_2, y]$ is isomorphic to $P_2 + P_2$, P_4 or K_3 with a pendant edge. In any case the subgraph $\Gamma[x, x_1, x_2, y]$ has rank 4, a contradiction.

For the assertion (ii), assume to the contrary, there exist $x_1 \in X, y_1 \in Y$ such that x_1 is not adjacent to y_1. Surely $x_1 \neq x$. As Γ has no isolated vertices, x_1 has a neighbor z, necessarily in Y by the result (i). The subgraph $\Gamma[x, x_1, y_1, z]$ has rank 4 whether y_1 and z are adjacent or not, also a contradiction. ■

Theorem 2.3. *Let Γ be a signed graph of order $n \geq 2$. Then $\eta(\Gamma) = n-2$ or equivalently $r(\Gamma) = 2$ if and only if Γ is a balanced complete bipartite graph together with some isolated vertices.*

Proof. The sufficiency is clear. So we only prove the necessity. First assume Γ has no isolated vertices. Choose an arbitrary vertex x in G. Let $Y = N(x), X = V(\Gamma) - Y$, both of which are nonempty. We assert any two vertices of Y are not adjacent; otherwise Γ contains a triangle which has rank 3 by Lemma 2.1. Then $\eta(\Gamma) \leq n-3$, a contradiction. Hence Γ is a complete bipartite graph by Lemma 2.2.

Next we show Γ is balanced. Otherwise Γ contains an unbalanced cycle, say C_{2k}. Further assume C_{2k} has the minimum length. By Lemma 2.1(iii), $k \geq 3$, as an unbalanced C_4 would have rank 4. Adding an edge between two vertices on C_{2k} with distance 3, the cycle is split into two cycles C_4 and C_{2k-2}, both being subgraphs of Γ. By the assumption on C_{2k}, C_4 is balanced, and hence C_{2k-2} is unbalanced, a contradiction.

Finally, if Γ contains isolated vertices, letting H be the subgraph of Γ by deleting those isolated vertices, then $r(H) = r(\Gamma) = 2$. So H is a balanced complete bipartite graph. The result now follows. ■

Let G be a signed graph and let v be a vertex of G. The *positive neighborhood* (respectively, *negative neighborhood*) of v is the set of vertices joining v with positive edges (respectively, negative edges).

Theorem 2.4. *Let Γ be a signed graph of order $n \geq 2$. Then $\eta(\Gamma) = n-3$ or equivalently $r(\Gamma) = 3$ if and only if Γ is a complete tripartite graph together with some isolated vertices, such that all vertices in each of three parts of the complete tripartite graph have the same positive neighborhoods and negative neighborhoods.*

Proof. If Γ is the graph as in the theorem, then Γ contains a triangle, and by Lemma 2.1, $r(\Gamma) \geq 3$. From the matrix of $A(\Gamma)$, all rows corresponding the vertices in a given part of the complete tripartite graph are same, which implies that $r(\Gamma) \leq 3$. So the sufficiency follows.

Now we prove the necessity. First assume Γ has no isolated vertices. Choose an arbitrary vertex x in G. Let $Y = N(x), X = V(\Gamma) - Y$, both of which are nonempty. By Lemma 2.2, any two vertices in X are not adjacent, and any vertex from X is adjacent to any vertex from Y.

We assert $r(\Gamma - X) = 2$. If $r(\Gamma - X) > 2$, then $r(G) = r(\Gamma - X) = 3$. So $\Gamma - X$ contains an induced subgraph of order 3 (i.e. a triangle) with rank 3. Therefore, G contains K_4 as a subgraph. If K_4 contains an unbalanced C_4, by Lemma 2.1(iii), $r(K_4) = 4$, a contradiction. Otherwise, by a little calculation we also get $r(K_4) = 4$, a contradiction. So $r(\Gamma - X) \leq 2$. If $r(\Gamma - X) < 2$, then $r(\Gamma - X) = 0$, and hence $\Gamma - X$ contains no edges. So, Γ is a complete bipartite graph, whose rank is an even number, a contradiction.

From above discussion, $\Gamma - X$ is a balanced complete bipartite graph together with some isolated vertices by Theorem 2.3. If $\Gamma - X$ contains isolated vertices, then Γ contains a triangle with an pendant edge as an induced subgraph. By Lemma 2.5 below, the latter subgraph has rank 4, a contradiction. So Γ is a complete tripartite graph. Write $A(\Gamma)$ as the following form:

$$A(\Gamma) = \begin{bmatrix} O & J_{12} & J_{13} \\ J_{12}^T & O & J_{23} \\ J_{13}^T & J_{23}^T & O \end{bmatrix} =: \begin{bmatrix} A_1 \\ A_2 \\ A_3 \end{bmatrix},$$

where each J_{ij} $(i \neq j)$ consists of entries 1 or -1. If there exists a row submatrix A_i, say A_1, has rank $r \geq 2$, then A_1^T also has rank r, which implies that $r(\Gamma) \geq 2r \geq 4$, a contradiction. So, each row of A_i is a copy of the first row of A_i for $i = 1, 2, 3$. The result follows in this case.

If Γ contains isolated vertices, letting H be the subgraph of Γ by deleting those isolated vertices, then $r(H) = r(\Gamma) = 3$. So H has the property as discussed above. The result follows. ■

At the end of this section, we introduce some graph transformations which preserve the nullity.

Lemma 2.5. [12] *Let Γ be a signed graph containing a pendant vertex, and let H be the induced subgraph of G obtained by deleting this pendant vertex together with the vertex adjacent to it. Then*

$$\eta(\Gamma) = \eta(H).$$

Corollary 2.6. *Let Γ be a signed graph of order $n \geq 4$ which is not a star. If Γ contains a pendant vertex, then $\eta(\Gamma) \leq n-4$.*

Proof. Let v be a pendant vertex of Γ, and let u be the vertex adjacent to v. By Lemma 2.5, $\eta(\Gamma) = \eta(\Gamma - v - u)$, and hence $r(\Gamma) = r(\Gamma - u - v) + 2$. As Γ is not a star, $\Gamma - u - v$ contains at least one edge so that $r(\Gamma - u - v) \geq 2$, which implies $r(\Gamma) \geq 4$, or equivalently $\eta(\Gamma) \leq n-4$. ■

Denote by $Pv_1v_2v_3$ a path on vertices v_1, v_2, v_3 with edges v_1v_2, v_2v_3. If a signed graph Γ contains a path $Pv_1v_2v_3$ which satisfies $d(v_2) = 2, v_1v_3 \notin E(\Gamma), N(v_1) \cap N(v_3) \setminus \{v_2\} = \emptyset$, then $Pv_1v_2v_3$ is called a *special path* of Γ.

Lemma 2.7. *Let Γ be a signed graph containing a special path $Pv_1v_2v_3$, where $\sigma(v_1v_2) = -1, \sigma(v_2v_3) = 1$. Suppose that $N(v_1) \setminus \{v_2\} \neq \emptyset$ and $v \in N(v_1) \setminus \{v_2\}$. Let Γ' be obtained from Γ by deleting the edge vv_1 and adding a new edge vv_3 with the same sign of vv_1. Then $\eta(\Gamma') = \eta(\Gamma)$.*

Proof. Without loss of generality, write $A(\Gamma)$ as follows, where the first to the fourth rows correspond to the vertices v_1, v_2, v_3, v respectively, and $H = \Gamma - \{v_1, v_2, v_3, v\}$.

$$A(\Gamma) = \begin{bmatrix} 0 & -1 & 0 & \sigma(vv_1) & \alpha \\ -1 & 0 & 1 & 0 & \mathbf{0} \\ 0 & 1 & 0 & 0 & \beta \\ \sigma(vv_1) & 0 & 0 & 0 & \gamma \\ \alpha^T & \mathbf{0}^T & \beta^T & \gamma^T & A(H) \end{bmatrix}$$

Taking the second row (column) to act on the fourth row (column) by elementary transformations, we arrive at a matrix:

$$M = \begin{bmatrix} 0 & -1 & 0 & 0 & \alpha \\ -1 & 0 & 1 & 0 & \mathbf{0} \\ 0 & 1 & 0 & \sigma(vv_1) & \beta \\ 0 & 0 & \sigma(vv_1) & 0 & \gamma \\ \alpha^T & \mathbf{0}^T & \beta^T & \gamma^T & A(H) \end{bmatrix}$$

Evidently M has the same rank as $A(\Gamma)$, is exactly the adjacency matrix of Γ'. The result follows. ■

Corollary 2.8. *Let Γ be a signed graph containing a special path $Pv_1v_2v_3$, where $\sigma(v_1v_2) = -1, \sigma(v_2v_3) = 1$. Let Γ' be a signed graph obtained from Γ by contracting the path into a single vertex. Then $\eta(\Gamma') = \eta(\Gamma)$.*

Proof. If $N(v_1) \setminus \{v_2\} \neq \emptyset$, deleting the edge vv_1 and adding a new edge vv_3 with same signs as vv_1 for each vertex $v \in N(v_1) \setminus \{v_2\}$, we will arrive at a graph H such that $\eta(H) = \eta(\Gamma)$ by Lemma 2.7. Noting that v_1 is a pendant vertex of H, by Lemma 2.5, $\eta(H) = \eta(H - v_1 - v_2)$. It is readily seen that $H - v_1 - v_2$ is isomorphic to Γ'. The result follow. ■

A *bicyclic graph G* is a simple connected graph such that $|E(G)| = |V(G)| + 1$.

Corollary 2.9. *Let B be a bicyclic signed graph of order n. If B contains a special path, then $\eta(B) \leq n-4$.*

Proof. Assume $Pv_1v_2v_3$ is a special path of B. There exists a sign function θ such that in the graph B^θ the sign of v_1v_2 is negative and the sign of v_2v_3 is positive. Noting that $\eta(B) = \eta(B^\theta)$, so we deal with the graph B^θ. By Corollary 2.8, we can get a new signed graph B' from B^θ such that $\eta(B') = \eta(B^\theta) = \eta(B)$ and $r(B) = r(B^\theta) = r(B') + 2$, where B' is obtained from B^θ by contracting the special path into a single vertex. As B is bicyclic, B' contains at least one edge, $r(B') \geq 2$ and hence $r(B) \geq 4$. The result follows. ■

3 Nullity of bicyclic signed graphs

There are two basic bicyclic graphs: ∞-graph and Θ-graph. An ∞-*graph*, denoted by $\infty(p,q,l)$, is obtained from two vertex-disjoint cycles C_p and C_q by connecting one vertex of C_p and one of C_q with a path P_l of length $l-1$ (in the case of $l=1$, identifying the above two vertices); and a Θ-*graph*, denoted by $\Theta(p,q,l)$, is a union of three internally disjoint paths $P_{p+1}, P_{q+1}, P_{l+1}$ of length p,q,l respectively with common end vertices, where $p,q,l \geq 1$ and at most one of them is 1. Observe that any bicyclic graph G is obtained from an ∞-graph or a Θ-graph (possibly) by attaching trees to some of its vertices.

Note that the simple bicyclic graphs (or balanced signed bicyclic graphs) of order n with nullity $n-3$ or $n-4$ have been characterized in [22]. In this section we will characterize the unbalanced signed bicyclic graphs of order n with nullity $n-3$ or $n-3$, and the signed bicyclic signed graphs (including simple bicyclic graphs) of order n with nullity $n-5$..

THEOREM 3.1. *Let B be an unbalanced bicyclic signed graph of order n. Then $\eta(B) \leq n-3$, with equality if and only if ${}^uB = \Theta(2,2,1)$ and the two triangles of B are both unbalanced.*

Proof. As B is unbalanced, by Theorem 2.3, $\eta(B) \leq n-3$. If $\eta(B) = n-3$, by Theorem 2.4, B is a complete tripartite graph, which implies ${}^uB = \Theta(2,2,1)$, and two triangles of B are both unbalanced. The sufficiency also follows from Theorem 2.4. ■



$G_1(n,p,q)$ $G_2(n,p,q)$

$G_3(n)$ $G_4(n)$

$G_5(n)$ $G_6(n)$

Fig. 3.1. Six bicyclic graphs with pendant vertices

THEOREM 3.2. *Let B be an unbalanced bicyclic signed graph of order n. Then $\eta(B) = n-4$ if and only if B is one of the following graphs with certain properties:*

(1) ${}^uB = G_1(n,p,q)$ *of Fig. 3.1, and B contains at least one unbalanced cycle,*

(2) ${}^uB = G_2(n,p,q)$ *of Fig. 3.1, and B contains at least one unbalanced cycle,*

(3) ${}^uB = G_3(n)$ *of Fig. 3.1, and the triangle of B is unbalanced and the quadrangle of B is balanced,*

(4) ${}^uB = G_4(n)$ *of Fig. 3.1, and the quadrangles of B containing the vertex v are unbalanced and the quadrangle of B containing no v is balanced,*

(5) ${}^uB = G_{12}$ *of Fig. 3.2, and B contains exactly one unbalanced cycle,*

(6) ${}^uB = G_{14}$ *of Fig. 3.3, and B contains an unbalanced triangle and a balanced quadrangle,*

(7) ${}^uB = G_{15}$ *of Fig. 3.3, and B has exactly one unbalanced triangle,*

(8) ${}^uB = G_{16}$ *of Fig. 3.3, and B^θ has an edge labeling as in Fig. 3.3 for some sign function θ.*

Proof. The sufficiency can be verified by Lemma 2.5, Lemma 2.1, Theorem 2.3, Corollary 2.8 or direct calculations.

Now assume $\eta(B) = n - 4$. We divide the discussion into cases.

Case 1: B contains pendant vertices. Let u be a pendant vertex of B, and let v be the vertex adjacent to u. By Lemma 2.5, $\eta(B) = \eta(B - u - v) = n - 4$. So $r(B - u - v) = 2$. By Theorem 2.3, $B - u - v$ is a balanced complete graph H together with some isolated vertices. As B is bicyclic, ${}^uH$ is a star, or C_4 or $\Theta(2, 2, 2)$. If ${}^uH = \Theta(2, 2, 2)$, then ${}^uB$ is $G_5(n)$ or $G_6(n)$ of Fig. 3.1. In this case B is also balanced as H is balanced. So this case cannot occur. If H is a star, then ${}^uB$ is $G_1(n, p, q)$ or $G_2(n, p, q)$ of Fig. 3.1, and B contains at least one unbalanced cycle. If $H = C_4$, then ${}^uB$ is $G_3(n)$ or $G_4(n)$ of Fig. 3.1. In addition, if ${}^uB = G_3(n)$, then the triangle is unbalanced and the quadrangle is balanced; if ${}^uB = G_4(n)$, then the quadrangle containing no v is balanced and the quadrangles containing v are unbalanced.

Case 2: B contains no pendant vertices. First assume that B is an ∞-graph and contains a special path P. Without loss of generality, assume P contains a positive edge and a negative edge. Now contracting the path P into a single vertex, by Corollary 2.8, the resulting graph, denoted by B', is still an ∞-graph, has the same nullity as B, and hence has rank 2. By Theorem 2.3, B' is a balanced complete bipartite graph, which is impossible. So B contains no special paths, and ${}^uB$ is one of graphs $G_i, i = 7, 8, \ldots, 12$, of Fig. 3.2. If ${}^uB$ is one of G_7, G_8, G_9, as B is unbalanced, B contains $C_3 + P_2$ or $C_4 + P_2$ as an induced subgraph, where C_3 or C_4 is unbalanced. By Lemma 2.1, $r(C_3 + P_2) = 5$, $r(C_4 + P_2) = 6$, a contradiction. So, it suffices to consider the case of ${}^uB$ being one of G_{10}, G_{11}, G_{12}. By a direct calculation, ${}^uB = G_{12}$, and B contains exactly one unbalanced cycle.


G_7 G_8 G_9

G_{10} G_{11} G_{12}

Fig. 3.2. Six ∞-graphs

Next assume that B is a Θ-graph. If B contains no special paths, then ${}^uB$ is one of the graphs G_{13}, G_{14}, G_{15} of Fig. 3.3. By a direct calculation, ${}^uB = G_{14}$ and B has an unbalanced triangle and a balanced quadrangle, or ${}^uB = G_{15}$ and B has exactly one unbalanced triangle. If B contains a special path P, without loss of generality, P starts at a vertex of degree 3, say $Pv_1v_2v_3$; see G_{16} of Fig. 3.3. Without loss of generality, the edge v_1v_2 is negative and v_2v_3 is positive. Now contracting the path P into a single vertex say u, by Corollary 2.8, the resulting graph $\bar{B}$ will have rank 2. By Theorem 2.3, $\bar{B}$ is a balanced complete bipartite graph, that is ${}^u\bar{B} = \Theta(2, 2, 2)$. By Theorem 1.1, there exists a sign function η defined on $V(\bar{B})$, such that all edges of $\bar{B}^\eta$ are positive. Returning to the original graph, ${}^uB = G_{16}$ of Fig. 3.3. Define a sign function θ on B such that $\theta(v_1) = \theta(v_2) = \theta(v_3) = \eta(u)$, and $\theta(v) = \eta(v)$ for any other vertex v. Then B^θ is the graph G_{16} with edge labeling as in Fig. 3.3. ■



G_{13} G_{14} G_{15} G_{16}

Fig. 3.3. Four Θ-graphs

Theorem 3.3. *Let B be a bicyclic signed graph of order n. Then $\eta(B) = n-5$ if and only if B is one of the following graphs with certain properties:*

(1) ${}^uB = G_{17}$ *of Fig. 3.4;*

(2) ${}^uB = G_{18}$ *of Fig. 3.4, and the two triangles of B have the same balanceness;*

(3) ${}^uB = G_{19}$ *of Fig. 3.4, and the two triangles of B have the same balanceness;*

(4) ${}^uB = G_{11}$ *of Fig. 3.2, and B is balanced;*

(5) ${}^uB = G_{12}$ *of Fig. 3.2, and the two triangles of B have the same balanceness;*

(6) ${}^uB = G_{14}$ *of Fig. 3.3, and the quadrangle of B is unbalanced;*

(7) ${}^uB = G_{20}$ *or* G_{21} *of Fig. 3.4,* B^θ *has an edge labeling as in Fig. 3.4 for some sign function θ.*

Proof. The sufficiency can be verified by Lemma 2.5, Lemma 2.1, Theorem 2.4, Corollary 2.8 or direct calculations.

Now assume that $\eta(B) = n-5$. We divide the discussion into cases.

Case 1: B contains pendant vertices. Let u be a pendant vertex of B, and let v be the vertex adjacent to u. By Lemma 2.5, $\eta(B) = \eta(B-u-v) = n-5$. So $r(B-u-v) = 3$. By Theorem 2.4, $B-u-v$ is a complete tripartite graph H together with some isolated vertices. As B is bicyclic, ${}^uH$ is a triangle, or $\Theta(2,2,1)$. Hence ${}^uB$ is one of the graphs G_{17}, G_{18}, G_{19} in Fig. 3.4, where the two triangles of B have the same balanceness for the last two cases by Theorem 2.4.

Case 2: B contains no pendant vertices. First assume that B is an ∞-graph and contains a special path P. Without loss of generality, assume P contains a positive edge and a negative edge. Now contracting the path P into a single vertex, by Corollary 2.8, the resulting graph, denoted by B', is still an ∞-graph, has the same nullity as B, and hence has rank 3. By Theorem 2.3, B' is a complete tripartite graph, which is impossible. So B contains no special paths, and ${}^uB$ is one of graphs $G_i, i = 7, 8, \ldots, 12$, of Fig. 3.2. By a little calculation, ${}^uB = G_{11}$ and B is balanced, or ${}^uB = G_{12}$ and two triangles of B have the same balanceness.

Next assume that B is a Θ-graph. If B contains no special paths, then ${}^uB$ is one of the graphs G_{13}, G_{14} of Fig. 3.3. By a direct calculation, ${}^uB = G_{14}$ and the quadrangle of B is unbalanced. If B contains a special path P, without loss of generality, P starts at a vertex of degree 3, say $Pv_1v_2v_3$; see G_{20}, G_{21} of Fig. 3.4. Without loss of generality, the edge v_1v_2 is negative and v_2v_3 is positive. Now contracting the path P into a single vertex say u, the resulting graph $\bar{B}$ will have rank 3. By Theorem 2.4, $\bar{B}$ is a complete tripartite graph, that is ${}^u\bar{B} = \Theta(2,2,1)$, where the two triangles of $\bar{B}$ have the same balanceness. By Theorem 1.1, there exists a sign function η defined on $V(\bar{B})$, such that all edges of $\bar{B}^\eta$ are positive or all edges of $\bar{B}^\eta$ are positive except the common edge of the two triangles. Returning to the original graph, ${}^uB = G_{20}$ or G_{21} of Fig. 3.4. Define a sign function θ on B such that $\theta(v_1) = \theta(v_2) = \theta(v_3) = \eta(u)$, and $\theta(v) = \eta(v)$ for any other vertex v. Then B^θ is the graph G_{20} or G_{21} with edge labeling as in Fig. 3.4.■

By Theorem 3.3, we easily get a characterization of simple bicyclic graphs (or balanced bicyclic signed graphs) of order n with nullity $n-5$.

Corollary 3.4. *Let B be a simple bicyclic graph of order n. Then $\eta(B) = n-5$ if and only if B is one of the following graphs: G_{11}, G_{12} of Fig. 3.2, G_{17}, G_{18}, G_{19} of Fig. 3.4.*



G_{17} G_{18} G_{19}

G_{20} G_{21}

Fig. 3.4. Five bicyclic graphs

References

[1] P. J. Cameron, J. Seidel, J. J. Tsaranov, Signed graphs, root lattices, and coxeter groups, *J. Algebra*, 164(1)(1994), 173-209.

[2] S. Chaiken, A combinatorial proof of the all minors matrix tree theorem, *SIAM J. Algebraic Discrete Methods*, 3(2)(1982), 319-329.

[3] B. Cheng, B. Liu, On the nullity of graphs, *Elec. J. Linear Algebra*, 16(2007), 60-67.

[4] G. J. Cheng, L.-H Huang, H.-G. Yeh, A characterization of graphs with rank 4, *Linear Algebra Appl.*, 434(8)(2011), 1793-1798.

[5] G. J. Cheng, L.-H Huang, H.-G. Yeh, A characterization of graphs with rank 5, *Linear Algebra Appl.*, 436(11)(2012), 4241-4250.

[6] L. Collatz, U. Sinogowitz, Spektren endlicher Grafen, *Abh. Math. Sem. Univ. Hamburg*, 21(1957), 63-77.

[7] D. Cvetković, M. Doob, H. Sachs, *Spectra of Graphs*, 3rd ed., Johann Ambrosius Barth, Heidelberg, 1995.

[8] B. Deradass, L. Archarya, Spectral criterion for cycle balance in networks, *J. Graph Theory*, 4(1)(1980), 1-11.

[9] S. Fallat, L. Hogben, The minimum rank of symmetric matrices described by a graph: a survey, *Linear Algebra Appl.*, 426(2007), 558-582.

[10] Y.-Z. Fan, Largest eigenvalue of a unicyclic mixed graph, *Appl. Math. J. Chinese Univ. Ser. B*, 19(2)(2004), 140-148.

[11] Y.-Z. Fan, K.-S. Qian, On the nullity of bipartite graphs, *Linear Algebra Appl.*, 430(2009), 2943-2949.

[12] Y.-Z. Fan, Y. Wang, Y. Wang, The nullity of unicyclic signed graphs, arXiv:1107.0400v1.

[13] S. Fiorini, I. Gutman, I. Sciriha, Trees with maximum nullity, *Linear Algebra and Appl.*, 397(2005), 245-251.

[14] I. Gutman, S.-L. Lee, J.-H. Sheu, C. Li, Predicting the nodal properties of molecular orbitals by means of signed graphs, *Bull. Inst. Chem. Academia Sinica*, 42(1995), 25-32.

[15] S.-C. Gong, Y.-Z. Fan, Z.-X. Yin, On the nullity of graphs with pendent trees, *Linear Algebra Appl.*, 433(7)(2010), 1374-1380.

[16] S.-C. Gong, G.-H. Xu, On the nullity of a graph with cut-points, *Linear Algebra Appl.*, 436(2012), 135-142.

[17] J.-M. Guo, W. Yan, Y.-N. Yeh, On the nullity and the matching number of unicyclic graphs, *Linear Algebra Appl.*, 431(8)(2009), 1293-1301.

[18] I. Gutman, I. Sciriha, On the nullity of line graphs of trees, *Discrete Math.*, 232(2001), 35-45.

[19] F. Harary, On the notion of balanced in a signed graph, *Michigan Math. J.*, 2(1)(1953), 143-146.

[20] Y. P. Hou, J. S. Li, On the Laplacian eigenvalues of signed graphs, *Linear Multilinear Algebra*, 51(1)(2003), 21-30.

[21] Y. P. Hou, Bounds for the least Laplacian eigenvalue of a signed graph, *Acta Math. Sinica, English Ser.*, 21(4)(2005), 955-960.

[22] S. Hu, X. Tan, B. Liu, On the nullity of bicyclic graphs, *Linear Algebra Appl.*, 429(7)(2008), 1387-1391.

[23] S.-L. Lee, C. Li, Chemical signed graph theory, *Int. J. Quantum Chem.*, 49(1994), 639-648.

[24] S.-L. Lee, R. R. Lucchese, Topological analysis of eigenvectors of the adjacency matrices in graph theory: the concept of internal connectivity, *Chem. Phys. Letters*, 137(3)(1987), 279-284.

[25] J. Li, A. Chang, W. C. Shiu, On the nullity of bicyclic graphs, *MATCH Commun. Math. Comput. Chem.*, 60 (2008), 21-36.

[26] S. Li, On the nullity of graphs with pendant vertices, *Linear Algebra Appl.*, 429(7)(2008), 1619-1628.

[27] W. Li, A. Chang, Describing the nonsingular unicyclic graph, *J. Math. Study*, 40(4)(2007), 442-445.

[28] H. C. Longuet-Higgins, Resonance structures and MO in unsaturated hydrocarbons, *J. Chem. Phys.*, 18(1950), 265-274.

[29] M. Nath, B. K. Sarma, On the null-spaces of acyclic and unicyclic singular graphs, *Linear Algebra Appl.*, 427(1)(2007), 42-54.

[30] F. S. Roberts, On balanced signed graphs and consistent marked graphs, *Elec. Notes Discrete Math.*, 2(1999), 94-105.

[31] P. K. Sahu, S.-L. Lee, Net-sign identity information index: a novel approach towards numerical characterization of chemical signed graph theory, *Chem. Phys. Letters*, 454(2008), 133-138.

[32] X. Tan, B. Liu, On the nullity of unicyclic graphs, *Linear Algebra Appl.*, 408(2005), 212-220.

[33] T. Zaslavsky, Signed graphs, *Discrete Appl. Math.*, 4(1)(1982), 47-74.

[34] W. Zhu, T.-Z. Wu, S.-B. Hu, A note on the nullity of unicyclic graphs, *J. Math. Res. Expo.*, 30(5)(2010), 817-824.